



CDC27

Catalog number

030073

Supplier

United States Biological

Cdc27 shares strong similarity with *Saccharomyces cerevisiae* protein Cdc27, and the gene product of *Schizosaccharomyces pombe* nuc 2. It is a component of the Anaphase Promoting Complex (APC), which is composed of eight protein subunits and is highly conserved in eucaryotic cells. The APC catalyzes the formation of the cyclin B ubiquitin conjugate that is responsible for the ubiquitin mediated proteolysis of B type cyclins. This protein and 3 other members of the APC complex contain the TPR (tetratricopeptide repeat), a protein domain important for protein protein interaction. This protein was shown to interact with mitotic checkpoint proteins including Mad2, p55CDC and BUBR1, and thus may be involved in controlling the timing of mitosis.

Applications

Western Blotting: 1:500-1:2000.

Immunohistochemistry: 1:200-1:1000.

ELISA: Suggested dilution 1:10000.

Optimal dilutions to be determined by the researcher.

Storage and Stability

May be stored at 4°C for short-term only. Aliquot to avoid repeated freezing and thawing. Store at -20°C. Aliquots are stable for 12 months. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap.

Immunogen

Purified recombinant fragment of human CDC27 expressed in *E. coli*.

Formulation

Ascites fluid, 0.03% sodium azide.

Purity

Ascites

Specificity

Human

Product Type

Mab

Source

human

Isotype

IgG1

**Grade**

Ascites

Applications

E IHC WB

Crossreactivity

Hu

Storage

-20°C

MW

91

Reference

1. Mol Cell Biol. 2004 May;24(9):3577-87.
2. J Biol Chem. 2004 Oct 1;279(40):42128-38.