



EphA5 (Ephrin type-A receptor 5, Tyrosine-protein kinase receptor EHK-1, Eph homology kinase-1, Receptor protein-tyrosine kinase HEK7, EPH-like kinase 7, EK7, EHK1, HEK7)

Catalog number

E3363-01G

Supplier

United States Biological

Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the γ phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains. The tyrosine kinase (TK) group is mainly involved in the regulation of cell-cell interactions such as differentiation, adhesion, motility and death. There are currently about 90 TK genes sequenced, 58 are of receptor protein TK (e.g. EGFR, EPH, FGFR, PDGFR, TRK, and VEGFR families), and 32 of cytosolic TK (e.g. ABL, FAK, JAK, and SRC families).

Applications

Suitable for use in ELISA and Western Blot. Other applications not tested.

Recommended Dilution

ELISA: 1:1,000

Western Blot: 1:50-1:100

Optimal dilutions to be determined by the researcher.

Storage and Stability

May be stored at 4°C for short-term only. Aliquot to avoid repeated freezing and thawing. Store at -20°C. Aliquots are stable for at least 12 months. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap.

Immunogen

Fusion recombinant protein of human EPHA5.

Formulation

Supplied as a liquid in PBS, 0.09% sodium azide.

Purity

Purified by ammonium sulfate precipitation.

Specificity

Recognizes human EPHA5.

**Product Type**

Pab

Source

human

Isotype

IgG

Grade

Purified

Applications

E WB

Crossreactivity

Hu

Storage

-20°C

MW

114.784

Reference

1. Fox, G.M., et al., Oncogene 10(5):897-905 (1995).